B.1 Incense Tree Aquilaria sinensis



Photo B.1.1.: General view of the transplanted individual AS-03.



Photo B.1.3.: General view of the transplanted individual AS-02.



Photo B.1.2.: Branch condition of the transplanted individual AS-03.



Photo B.1.4.: Branch condition of the transplanted individual AS-02.



B.2 Lamb of Tartary *Cibotium barometz*



Photo B.2.1.: General view of the transplanted individual CB-01.



Photo B.2.3.: Leaf condition of the transplanted individual CB-01.



Photo B.2.2.: Leaf condition of the transplanted individual CB-01.



Photo B.2.4.: Leaf condition of the transplanted individual CB-01.



Bottlebrush Orchid Goodyera procera *B.3*



Photo B.3.1: Individual GP-01. Partially wilted leaf.





Photo B.3.3: Individual GP-02.



Photo B.3.4: Individual GP-03. Perforated leaves.





Photo B.3.5: Individual GP-03. Perforated leaves.



Photo B.3.7: Individual GP-04. Wilted leaf.



Photo B.3.6: Individual GP-04. Wilted leaf.



Photo B.3.8: Individual GP-05. Partially wilted leaf.





Photo B.3.9: Individual GP-06. Partially wilted leaf.



Photo B.3.10: Individual GP-06. Partially chlorotic and wilted leaf.



Photo B.3.11: Individual GP-07.




Photo B.3.12: Individual GP-07. Single chlorotic leaf.





Photo B.3.13: Individual GP-08. Chlorotic leaves.



Photo B.3.15: Individual GP-09.




Photo B.3.14: Individual GP-08. Minor chlorotic leaves.



Photo B.3.16: Individual GP-10. Holes in leaves.





Photo B.3.17: Individual GP-11. Partially wilted leaves.



Photo B.3.18: Individual GP-12. Perforated leaves.



Photo B.3.19: Individual GP-13.



Photo B.3.20: Individual GP-14. Chlorotic leaves.





Photo B.3.21: Individual GP-15. Partially chlorotic leaves.



Photo B.3.22: Individual GP-16.



Photo B.3.23: Individual GP-17. Partially wilted leaf.



Photo B.3.24: Individual GP-18. Partially wilted leaves.





Photo B.3.25: Individual GP-19.



Photo B.3.26: Individual GP-19. Chlorotic and partially wilted leaves.

